arXiv:1704.04443v1 [math.FA] 13 Apr 2017

ON n-TUPLET FIXED POINTS FOR NONCOMPACT MULTIVALUED MAPPINGS VIA MEASURE OF NONCOMPACTNESS

DERYA SEKMAN, NOUR EL HOUDA BOUZARA, AND VATAN KARAKAYA

Abstract. In this work, by using measure of noncompactness, some results on the existence of n-tuplet fixed points for multivalued set contraction mappings are proved. As an application, the existence of solution for a system of integral inclusions is studied.

1. Introduction

Fixed point theory is an important working area in mathematics which is closely related to real world problems and has many applications different fields of science. It has played an important role in solving problems of uniqueness and existence of the nonlinear works as well as analysis, topology and geometry. In the literature, it has applications in many sciences like engineering [1], economy [2], optimization [3], game theory [4] and medicine [5]. Fixed point theory was originated by Poincaré in 1912. Brouwer [6] given fixed point theorem for n-dimensional space. Banach [7] proved Banach contraction principle which supplied find fixed point theorem using contraction mapping. Later, Schauder [8] proved an extension of the Brouwer's fixed point theorem to spaces of infinite dimension under a condition of compactness.

In the fixed point theory, it can be seen to development in different directions. One of them is single valued mappings which are many works. One can find the following references [9, 10]. However, we can do more important studies with multivalued mappings compared to single valued mappings since they have a direct application to the real world [11, 12]. According to development of literature in this area, Kakutani [13] extended Brouwer's fixed point theorem to multivalued mappings. Afterward, Nadler [14] extended the Banach contraction principle from single valued mapping to multivalued mapping and proved a fixed point theorem using Hausdorff metric for the multivalued contraction. Later, Nadler's fixed point theorem was extended and generalized for class of various multivalued mappings by many authors [15, 16].

Measure of noncompactness has played a fundamental role in the study of single valued and multivalued mappings, especially metric and topological fixed point theory. It is very useful tool due to compactness condition guarantees the existence of fixed point. The measure of noncompactness was defined and studied by Kuratowski [17] and Hausdorff [18]. Darbo [19] used this measure to generalize both the

1991 *Mathematics Subject Classification.* 47H9, 47H10, 34A60.

Key words and phrases. Measure of noncompactness, n-tuplet fixed point, multivalued set contraction mapping, system of differential inclusions.

Schauder's fixed point theorem and the Banach's contraction principle for condensing operators. Recently, measure of noncompactness have been used in differential equations, integral equations, nonlinear equations etc., some of works are noted in [20, 21].

The order and partially ordered metric spaces are very important in the works on fixed point theory. By using two basic concepts, Guo and Lakshmikantham [22] first gave some existence theorems of the coupled fixed point for both continuous and discontinuous operators, and then they offered some applications to the initial value problems of ordinary differential equations with discontinuous right-hand sides. Bhaskar and Lakshmikantham [23] introduced a coupled fixed point and established some coupled fixed point theorems for partially ordered metric space. By the similar idea, Berinde and Borcut [24] introduced a tripled fixed point theorem for nonlinear mapping in partially ordered complete metric space. Ertürk and Karakaya [25] introduced the concept of n-tuplet fixed point theorem and studied existence and uniqueness of fixed point of contractive type mappings in partially ordered metric spaces. Moreover, by using the condensing operators, Aghajani et al. [26] presented some results on the existence of coupled fixed point and also Karakaya et al. [27] gave some results concerning the existence of tripled fixed point via measure of noncompactness.

The existence of fixed point for various contractive mappings has been studied by many authors under different conditions. The concept of coupled fixed point for multivalued mappings was introduced by Samet and Vetro [28] and they presented coupled fixed point theorem for multivalued nonlinear contraction mappings in partially ordered metric space. Rao et al. [29] obtained a tripled coincidence fixed point theorems for multivalued mappings in partially ordered metric space.

In this paper, by using condensing operator, we investigate n-tuplet of fixed point theorem of multivalued mapping on Banach space. Finally, we will give an application including the system of integral inclusion and also give some special cases as coupled and tripled fixed point.

2. Preliminaries

Throughout this paper, let E be a Banach space and $P(E)$ (also noted 2^E) be the set of all subsets of E. In further we denote

$$P_k(E) = \{X \subset E,\ X \text{ is nonempty and has a property } k\},$$

where $P_{rcp}(E), P_{cl,bd}(E), P_{cl,cv}(E)$ denote the classes of all relatively compact, closed-bounded and closed-convex subsets of E respectively.

A mapping $T : E \to P_k(E)$ is called a multivalued mapping or set valued mapping on E into E. A point $x \in E$ is called a fixed point of T if $x \in Tx$.

Definition 1 (see; [30]). *A mapping $\mu : P_{cl,bd}(X) \to \mathbb{R}^+$ is called a measure of noncompactness if it satisfied the following conditions:*

(M_1) $\varnothing \neq \mu^{-1}(0) \subset P_{rcp}(X)$,
(M_2) $\mu(\bar{A}) = \mu(A)$, where $\bar{A}$ denotes the closure of A,
(M_3) $\mu(conv\ A) = \mu(A)$, where $conv\ A$ denotes the convex hull of A,
(M_4) μ is nondecreasing,

(M_5) If $\{A_n\}$ is a decreasing sequence of sets in $P_{cl,bd}(X)$ satisfying $\lim_{n\to\infty}\mu(A_n)=0$, then the intersection

$$A_\infty=\bigcap_{n=1}^{\infty}A_n$$

is nonempty.

If (M_4) holds, then $A_\infty\in P_{rcp}(X)$. For this, let $\lim_{n\to\infty}\mu(A_n)=0$. As $A_\infty\subseteq A_n$ for each $n=0,1,2,...$; by the monotonicity of μ, we obtain

$$\mu(A_\infty)\leq\lim_{n\to\infty}\mu(A_n)=0.$$

So, by (M_1), we get that A_∞ is nonempty and $A_\infty\in P_{rcp}(X)$.

Theorem 1 (see; [31]). *Let X be a closed and convex subset of a Banach space E. Then every compact, continuous map $T:X\to X$ has at least one fixed point.*

Theorem 2 (see; [19]). *Let X be a nonempty, bounded, closed and convex subset of a Banach space E and let $T:X\to X$ be a continuous mapping. Suppose that there exists a constant $k\in[0,1)$ such that*

$$\mu(T(X))\leq k\mu(X)$$

for any subset X of E, then T has a fixed point.

Definition 2 (see; [30]). *A multivalued mapping $T:E\to P_{cl,bd}(E)$ is called D-set-Lipschitz if there exists a continuous nondecreasing function $\varphi:\mathbb{R}^+\to\mathbb{R}^+$ such that*

$$\mu(T(X))\leq\varphi(\mu(X))$$

for all $X\in P_{cl,bd}(E)$ with $T(X)\in P_{cl,bd}(E)$, where $\varphi(0)=0$. Generally, we call the function φ to be a D-function of T on E.

When $\varphi(r)=kr$, $k>0$, T is called a k-set-Lipschitz mapping and if $k<1$, then T is called a k-set-contraction on E.

If $\varphi(r)<r$ for $r>0$, then T is called a *nonlinear D-set-contraction* on E.

Lemma 1 (see; [32]). *If φ is a D-function with $\varphi(r)<r$ for $r>0$, then*

$$\lim_{n\to\infty}\varphi^n(t)=0$$

for all $t\in[0,\infty)$.

Theorem 3 (see; [30]). *Let X be a nonempty, bounded, closed and convex subset of a Banach space E and let $T:X\to P_{cl,cv}(X)$ be a closed and nonlinear D-set-contraction. Then T has a fixed point.*

As a consequence of Theorem 3, we obtain a fixed point theorem of Darbo ([19]) type for linear set-contractions.

Corollary 1 (see; [30]). *Let X be a bounded, closed and convex subset of a Banach space E and let $T:X\to P_{cl,cv}(X)$ be a closed and k-set-contraction. Then T has a fixed point.*

Definition 3 (see; [33]). *Let X be a topological space, 2^X be the family of all subsets of X and T be a mapping of X into 2^X such that Tx is nonempty, for all $x\in X$. Then the mapping T is called upper semicontinuous if for each closed subset C of X,*

$$T^{-1}(C)=\{x\in X:Tx\cap C\neq\varnothing\}$$

is closed.

Definition 4 (see; [30]). *A mapping $\mu : P_k(E) \to \mathbb{R}^+$ is called nondecreasing if $A, B \in P_k(E)$ are any two sets with $A \subseteq B$, then $\mu(A) \leq \mu(B)$, where $\subseteq$ is an order relation by inclusion in $P_k(E)$.*

Lemma 2 (see; [34]). *Let X be a Banach space and F be a Caratheodory multivalued mapping. Let $\Phi : L^1(H;X) \to C(H;X)$ be linear continuous mapping. Then,*

$$\begin{aligned} \Phi \circ S_F : C(H;X) &\to \mathcal{P}_{cl;cv}(C(H;X)) \\ u &\to (\Phi \circ S_F)\, u := \Phi(S_F(u)), \end{aligned}$$

is a closed graph operator in $C(H;X) \times C(H;X)$.

Lemma 3 (see; [35]).

1. Let $A \subseteq C(H;X)$ is bounded, then $\mu(A(t)) \leqslant \mu(A)$ for all $t \in H$, where $A(t) = \{y(t), y \in A\} \subset X$. Furthermore, if A is equicontinuous on H, then $\mu(A(t))$ is continuous on H and $\mu(A) = \sup\{\mu(A(t)), t \in H\}$.
2. If $A \subset C(H;X)$ is bounded and equicontinuous, then

$$\mu\left(\int_0^t A(s)\, ds\right) \leqslant \int_0^t \mu(A(s))\, ds,$$

for all $t \in H$, where $\int_0^t A(s)\, ds = \left\{\int_0^t x(s)\, ds : x \in A\right\}$.

3. n-tuplet Fixed Point Theorems of Multivalued Mappings and Some Related Results

In this section, we investigate n-tuplet fixed point property for multivalued mapping. Afterward, we will give some applications for special cases $n = 2$, that is, coupled fixed point.

Definition 5. *Let X be a nonempty set and $G : X^n \to P(X)$ be a given mapping. An element $(x_1, x_2, x_3, ..., x_n) \in X^n$ is called an n-tuplet fixed point of G if*

$$\begin{aligned} x_1 &\in G(x_1, x_2, x_3, ..., x_n) \\ x_2 &\in G(x_2, x_3, ..., x_n, x_1) \\ &\vdots \\ x_n &\in G(x_n, x_1, x_2, ..., x_{n-1}). \end{aligned}$$

Remark 1. *If we take as special cases $n = 2$ and $n = 3$ in the Definition 5, respectively, we get coupled fixed point (see; [28]) and tripled fixed point (see; [29]).*

Theorem 4. *(see; [36]) Let $\mu_1, \mu_2 ..., \mu_n$ are measures of noncompactness in Banach spaces $E_1, E_2 ..., E_n$ respectively. Suppose that the function $F : [0, \infty)^n \to [0, \infty)$ is convex and $F(x_1, x_2, ..., x_n) = 0$ if and only if $x_i = 0$ for $i = 1, 2, ..., n$. Then*

$$\widetilde{\mu}(X) = F(\mu_1(X_1), \mu_2(X_2) ..., \mu_n(X_n)),$$

defines a measure of noncompactness in $E_1 \times E_2 \times ... \times E_n$ where X_i denotes the natural projection of X into E_i, for $i = 1, 2, ..., n$.

Remark 2. *We can notice that by taking*

$$F(x_1, x_2, x_3, ..., x_n) = \max\{x_1, x_2, x_3, ..., x_n\},$$

or

$$F(x_1, x_2, x_3, ..., x_n) = x_1 + x_2 + x_3 + ... + x_n,$$

for any $(x_1, x_2, x_3, ..., x_n) \in [0, \infty)^n$*, conditions of Theorem 4 are satisfied. Therefore,*

$$\widetilde{\mu}(X) := \max(\mu(X_1), \mu(X_2), ..., \mu(X_n)),$$

or

$$\widetilde{\mu}(X) := \mu(X_1) + \mu(X_2) + \cdots + \mu(X_n),$$

defines measures of noncompactness in the space E^n*, where* X_i*,* $i = 1, 2, ..., n$ *are the natural projections of* X *on* E_i*.*

After this step, we will give an important theorem for the fixed point of multivalued mappings under measure of noncompactness condition.

Theorem 5. *Let* X *be a nonempty, bounded, closed and convex subset of a Banach space* E *and let* μ *be an arbitrary measure of noncompactness in it. Let* $\varphi : \mathbb{R}^+ \to \mathbb{R}^+$ *be a nondecreasing and upper semicontinuous function such that* $\varphi(r) < r$ *for all* $r > 0$*. Suppose that the* $G : X_1 \times X_2 \times \cdots \times X_n \to P_{cl,cv}(X)$ *is continuous multivalued operator satisfying*

$$\mu(G(X_1 \times X_2 \times \cdots \times X_n)) \leqslant \varphi\left(\frac{\mu(X_1) + \mu(X_2) + \cdots + \mu(X_n)}{n}\right)$$

for all $X_1, X_2, ..., X_n \subset X$*. Then* G *has at least an* n*-tuplet fixed point.*

Proof. By considering Remark 2, we define the measure of noncompactness $\tilde{\mu}$ by

$$\widetilde{\mu}(X) := \mu(X_1) + \mu(X_2) + \cdots + \mu(X_n)$$

and also the mapping $\tilde{G}(X) := G(X_1 \times X_2 \times \cdots \times X_n)$. We prove that $\widetilde{G}$ satisfies all the conditions of Theorem 3. Then, we obtain

$$\begin{aligned}
\widetilde{\mu}\left(\widetilde{G}(X)\right) &= \widetilde{\mu}(G(X_1 \times X_2 \times \cdots \times X_n)) \\
&= \widetilde{\mu}(G(x_1, x_2, x_3, ..., x_n), G(x_2, x_3, ..., x_n, x_1), ..., G(x_n, x_1, x_2, ..., x_{n-1})) \\
&= \mu(G(x_1, x_2, x_3, ..., x_n)) + \mu(G(x_2, x_3, ..., x_n, x_1)) + \cdots + \mu(G(x_n, x_1, x_2, ..., x_{n-1})) \\
&\leqslant \varphi\left(\frac{\mu(X_1) + \mu(X_2) + \cdots + \mu(X_n)}{n}\right) + \varphi\left(\frac{\mu(X_2) + \mu(X_3) + \cdots + \mu(X_1)}{n}\right) \\
&\quad + \cdots + \varphi\left(\frac{\mu(X_n) + \mu(X_1) + \cdots + \mu(X_{n-1})}{n}\right) \\
&= n\varphi\left(\frac{\mu(X_1) + \mu(X_2) + \cdots + \mu(X_n)}{n}\right).
\end{aligned}$$

Now,

$$\frac{1}{n}\widetilde{\mu}\left(\widetilde{G}(X)\right) \leq \varphi\left(\frac{\mu(X_1) + \mu(X_2) + \cdots + \mu(X_n)}{n}\right)$$

and taking $\widetilde{\mu}' = \frac{1}{n}\widetilde{\mu}$, we get

$$\widetilde{\mu}'\left(\widetilde{G}(X)\right) \leqslant \varphi\left(\widetilde{\mu}'(X)\right).$$

Also, $\widetilde{\mu}'$ is a measure of noncompactness. Then, by Theorem 3 we obtain that G has at least an n-tuplet fixed point. □

Remark 3. *If we take μ measure of noncompactness in Theorem 5 by*

$$\widetilde{\mu}(X) := \max(\mu(X_1), \mu(X_2), ..., \mu(X_n)),$$

we can obtained same results.

4. Applications To Inclusions Systems

The multivalued fixed point theorems of this paper can have some nice applications to differential and integral systems of inclusions as an example we study the solvability of a system of differential inclusions.

Let the following differential system

$$\begin{cases} x'(t) \in A(t)x(t) + G(t, x(t), y(t)), & t \in [0,b] \\ y'(t) \in A(t)y(t) + F(t, y(t), x(t)), & t \in [0,b] \end{cases} \tag{4.1}$$

with

$$x(0) = \varphi(x,y),\ y(0) = \varphi(y,x) \tag{4.2}$$

where G is an upper Caratheodory multimap, $\varphi : C([0,b], X) \to X$ is a given multivalued function. $\{A(t) : t \in [0,b]\}$ is a family of linear closed unbounded operators on X with domain $D(A(t))$ independent of t that generate an evolution system of operators $\{U(t,s) : t,s \in \Delta\}$ with $\Delta = \{(t,s) \in [0,b] \times [0,b] : 0 \leqslant s \leqslant t \leqslant b\}$.

Define the set

$$S_G(x,y) = \left\{ g \in L^1([0,b], X) : g(t) \in G(t, x(t), y(t)) \right\}.$$

Definition 6. *A family $\{U(t,s)\}_{(t,s)\in\Delta}$ of bounded linear operators $U(t,s) : X \to X$ where $(t,s) \in \Delta := \{(t,s) \in J \times J : 0 \leqslant s \leqslant t < +\infty\}$ for $J = [0,b]$ is called an evolution system if the following properties are satisfied,*

1. *$U(t,t) = I$ where I is the identity operator in X and $U(t,s)\ U(s,\tau) = U(t,\tau)$ for $0 \leqslant \tau \leqslant s \leqslant t < +\infty$,*
2. *The mapping $(t,s) \to U(t,s)\ y$ is strongly continuous, that is, there exists a constant $M > 0$ such that*

$$\|U(t,s)\| \leqslant M \quad \text{for any } (t,s) \in \Delta.$$

An evolution system $U(t,s)$ is said to be compact if $U(t,s)$ is compact for any t-$s > 0$. $U(t,s)$ is said to be equicontinuous if $\{U(t,s)x : x \in M\}$ is equicontinuous at $0 \leqslant s < t \leqslant b$ for any bounded subset $B \subset X$. Clearly, if $U(t,s)$ is a compact evolution system, it must be equicontinuous. The inverse is not necessarily true.

More details on evolution systems and their properties could be found on the books of Ahmed [37], Engel and Nagel [38] and Pazy [39].

Definition 7. *We say that the couple $(x(t), y(t)) \in C([0,b], X) \times C([0,b], X)$ is a mild solution of the evolution system (4.1) − (4.2) if it satisfies the following integral system*

$$\begin{cases} x(t) = U(t,0)\ \varphi(x,y) + \int_0^t U(t,s)\ g(s)\ ds & \text{for } g \in S_G(x,y) \\ y(t) = U(t,0)\ \varphi(y,x) + \int_0^t U(t,s)\ g(s)\ ds & \text{for } g \in S_G(y,x) \end{cases}, \tag{4.3}$$

for all $t \in [0,b]$.

Theorem 6. *Assume the following hypotheses*

$(H1)$ $\{A(t) : t \in J\}$ is a family of linear operators. $A(t) : D(A) \subset X \to X$ generates an equicontinuous evolution system $\{U(t,s) : (t,s) \in \Delta\}$ and

$$|U(t,s)| \leqslant M.$$

$(H2)$ The multifunction $G : J \times C([0,b] \times X \times X) \longrightarrow \mathcal{P}_{cl,cv}(X)$ is an upper Carathéodory with respect to x and y and $\varphi : C(J;X) \to X$ is compact and

$$\mu(G(t, W \times W)) < k\mu\left(\frac{W \times W}{2}\right) \text{ for any } t \in J.$$

$(H3)$ There exists a constant $r > 0$ such that

$$M\left[\|\varphi(x,y)\| + \{\|g(t)\|_1 : g \in S_G(x,y), x \in A_0\}\right] \leqslant r$$

and

$$M\left[\|\varphi(y,x)\| + \{\|g(t)\|_1 : g \in S_G(y,x), y \in A_0\}\right] \leqslant r$$

where, $A_0 = \{z \in C(J;X) : \|z(t)\| \leqslant r \text{ for all } t \in J\}$ hold, then the non local system $(4.1)-(4.2)$ has at least one mild solution in the space $C(J,X)$.

Proof. To solve problem $(4.1)-(4.2)$ we transform it to the following fixed point problem.

Consider the multivalued operator $N : C([0,b];X;X) \to \mathcal{P}(C([0,b];X))$ defined by,

$$N(x,y) = \left\{h \in C(J;X) : h(t) = U(t,0)\varphi(x,y) + \int_0^t U(t,s)\ g(s)\ ds, \text{ with } g \in S_G(x,y)\right\}.$$

Clearly coupled fixed point of the operator N are mild solutions of system $(4.3)-(4.2)$.

Obviously for each $y \in C([0,b];X)$, the set $S_G(x,y)$ is nonempty since, by $(H2)$, G has a measurable selection (see [40]).

Let show that N has a coupled fixed point. For that, we need to satisfy all the conditions Theorem 5.

Let $A_0 = \{z \in C([0,b];X) : \|z(t)\| \leqslant r \text{ for all } t \in [0,b]\}$. We can easily notice that A_0 is closed, bounded and convex.

To show that $N(A_0 \times A_0) \subseteq A_0$, we need first to prove that the family

$$\left\{\int_0^t U(t,s)\ f(s)\ ds\ : f \in S_F(y) \text{ and } y \in A_0\right\}$$

is equicontinuous for $t \in J$, that is, all the functions are continuous and they have equal variation over a given neighbourhood.

In view of $(H1)$ we have that funtions in the set $\{U(t,s) : (t,s) \in \Delta\}$ are equicontinuous, i.e, for every $\varepsilon > 0$ there exists $\delta > 0$ such that $|t-\tau| < \delta$ implies $\|U(t,s) - u(\tau,s)\| < \varepsilon$ for all $U(t,s) \in \{U(t,s) : (t,s) \in \Delta\}$

Then, given some $\varepsilon > 0$ let $\delta = \frac{\varepsilon'}{\varepsilon\|g\|_\infty}$ such that $|t-\tau| < \delta$, we have

$$\left|\int_0^t U(t,s)\ g(s)\ ds - \int_0^\tau U(\tau,s)gf(s)\ ds\right| \leqslant \int_\tau^t |U(t,s) - U(\tau,s)|\,|g(s)|\,ds.$$

Regarding the fact that $\{U(t,s) : (t,s) \in \Delta\}$ is equicontinuous then

$$\begin{aligned}\left|\int_0^t U(t,s)\ g(s)\ ds - \int_0^\tau U(\tau,s)\ g(s)\ ds\right| &\leqslant \varepsilon \|g\|_\infty |t-\tau| \\ &< \varepsilon \|g\|_\infty \frac{\varepsilon'}{\varepsilon \|g\|_\infty} = \varepsilon'.\end{aligned}$$

Hence we conclude that $\left\{\int_0^t U(t,s)\ g(s)\ ds\ : g \in S_G(x,y) \text{ and } (x,y) \in A_0 \times A_0\right\}$ is equicontinuous for $t \in J$.

Now, let show that $N(A_0 \times A_0) \subseteq A_0$. Let for $t \in J$,

$$\begin{aligned}|h(t)| &= \left|U(t,0)\,\varphi(x,y) + \int_0^t U(t,s)\ g(s)\ ds\right| \\ &\leqslant |U(t,0)\,\varphi(x,y)| + \int_0^t |U(t,s)\ g(s)|\, ds \\ &\leqslant M\|\varphi(x,y)\| + M\|g\|_1 \\ &= M\left[\|\varphi(x,y)\| + \|g\|_1\right] \leqslant r,\end{aligned}$$

thus $N(A_0 \times A_0) \subseteq A_0$.

In further, it is easy to see that N has convex valued.

Now, let show that N has a closed graph, let $x_n \to x$, $y_n \to y$ and $h_n \to h$ such that $h_n(t) \in N(x_n, y_n)$ and let show that $h(t) \in N(x,y)$.

Then, there exists a sequence $g_n \in S_G(x_n, y_n)$ such that

$$h_n(t) = U(t,0)\,\varphi(x_n,y_n) + \int_0^t U(t,s) g_n(s) ds.$$

Consider the linear operator $\Phi : L^1([0,b];X) \to C([0,b];X)$ defined by

$$\Phi f(t) = \int_0^t U(t,s) g_n(s) ds.$$

Clearly, Φ is linear and continuous. Then from Lemma 2, we get that $\Phi \circ S_G(x,y)$ is a closed graph operator. In further, we have

$$h_n(t) - U(t,0)\,\varphi(x_n,y_n) \in \Phi \circ S_G(x,y).$$

Since $x_n \to x$, $y_n \to y$ and $h_n \to h$, then

$$h(t) - U(t,0)\,\varphi(x,y) \in \Phi \circ S_G(x,y).$$

That is, there exists a function $g \in S_G(x,y)$ such that

$$h(t) = U(t,0)\,\varphi(y) + \int_0^t U(t,s) g(s) ds.$$

Therefore N has a closed graph, hence N has closed values on $C([0,b] \times X \times X, X)$.

We know that the family $\left\{\int_0^t U(t,s) f(s) ds, f \in S_F(W(t))\right\}$ is equicontinuous, hence by Lemma 3, we have

$$\begin{aligned}\mu\left(\int_0^t U(t,s)g(s)ds,\ g \in S_G(W(t) \times W(t))\right) &\leqslant \int_0^t \mu\left(U(t,s)g(s),\ g \in S_G(W(t) \times W(t))\right) ds \\ &\leqslant M\int_0^t \mu\left(g(s),\ g \in S_G(W(t) \times W(t))\right) ds \\ &\leqslant Mt\mu\left(G(t, W(t))\right).\end{aligned}$$

Therefore

$$\begin{aligned}\mu\left(N\left(W\times W\right)\right) &= \mu N\left(U\left(t,0\right)\varphi(W\left(t\right)\times W\left(t\right))+\int_0^t U(t,s)g(s)ds, g\in S_G\left(W\left(t\right)\times W\left(t\right)\right)\right)\\ &\leqslant \mu\left(U\left(t,0\right)\varphi(W\left(t\right)\times W\left(t\right))\right)+\mu\left(\int_0^t U(t,s)g(s)ds, g\in S_G\left(W\left(t\right)\times W\left(t\right)\right)\right)\\ &\leqslant M\mu\left(\varphi(W\left(t\right)\times W\left(t\right))\right)+Mt\mu\left(G\left(t,W\left(t\right)\times W\left(t\right)\right)\right).\end{aligned}$$

In view of $(H2)$, we get

$$\mu\left(N\left(W\times W\right)\right)\leqslant Mbk\mu\left(\frac{W\times W}{2}\right).$$

Therefore, for $Mbk<1$ we obtained that N has at least one coupled fixed point. Hence, the system $(4.1)-(4.2)$ has at least one solution.

References

[1] T. Fleiner, A fixed-point approach to stable matchings and some applications, Mathematics of Operations Research. 28 (2003), 103-126.

[2] E. A. Ok, Real analysis with economic applications, Princeton University, Princeton, (2007).

[3] L. C. Ceng, Q. Ansari, J. C. Yao,Some iterative methods for finding fixed points and for solving constrained convex minimization problems, Nonlinear Analysis: Theory, Methods & Applications. 74 (2011) ,5286-5302.

[4] K. C. Border, Fixed point theorems with applications to economics and game theory, Cambridge University, Cambridge, (1989).

[5] M. Chen, W. Lu, Q. Chen, K. J. Ruchala, G.H. Olivera, A simple fixed-point approach to invert a deformation field, Medical Physics. 35 (2008), 81-88.

[6] L. E. J. Brouwer, Uber abbilldung von mannigfaltigkeiten, Math. Ann. 71 (1912), 97-115.

[7] S. Banach, Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales, Fund. Math. 3 (1922), 133-181.

[8] J. Schauder, Der fixpunktsatzin funktionalraiimen, Studia Mathematika. 2 (1930), 171-180.

[9] L. P. Belluce, W. A. Kirk, Fixed point theorems for families of contraction mappings, Pasific J. Math. 18 (1966), 213-217.

[10] L. P. Belluce, W. A. Kirk, Nonexpansive mappings and fixed points in Banach spaces, Illinois J. Math. 11 (1967), 474-479.

[11] J. F. Nash, Equilibrium points in n-person games, Proc. Nat. Acad. Sci. 36 (1950), 48-49.

[12] A. Petruşel, Operatorial inclusions, House of the Book of Science, Cluj Napoka, (2002).

[13] S. Kakutani, A generalization of Brouwer's fixed point theorem, Duke Math. 8 (1941), 457-459.

[14] S. B. Nadler, Multivalued contraction mappings, Path. J. Math. 30 (1969), 475-488.

[15] S. Reich, Fixed points of contractive functions, Boll Un Mat Ital. 5 (1972), 26-42.

[16] N. Mizoguchi, W. Takahashi, Fixed points theorems for multivalued mappings on complete metric spaces, J. Math. Anal. Appl. 141 (1989), 177-188.

[17] K. Kuratowski, Sur les escapes complets, Fund. Math. 5 (1930), 301-309.

[18] L. S. Goldenštein, I. T. Gohberg, A. S. Markus, Investigatons of some properties of bounded linear operators with their q-norms, Ŭ cen. Zap. Kishinevsk. 29 (1957), 29-36.

[19] J. Banas, K. Goebel, Measures of noncompactness in Banach spaces, Lecture Notes in Pure and Applied Mathematics, 60 (1980), Dekker, New York

[20] A. Aghajani, J. Banas, Y. Jalilian, Existence of solutions for a class of nonlinear volterra singular integral equations, Comput. Math. Appl. 62 (2011), 1215-1227.

[21] J. Banaś, Measure of noncompactness in the spaces of continuous tempered functions, Demonstratio Math. 14 (1981), 127-133.

[22] D. Guo, V. Lakshmikantham, Coupled fixed points of nonlinear operators with applications, Nonlinear Anal. Theory Methods Appl. 11 (1987), 623-632.

[23] T. G. Bhaskar, V. Lakshmikantham, Fixed point theorems in partially ordered metric spaces and applications, Nonlinear Analysis. 65 (2006), 1379-1393.

[24] V. Berinde, M. Borcut, Tripled fixed point theorems for contractive type mappings in partially ordered metric spaces, Nonlinear Anal. Theory Methods Appl. 74 (2011), 4889-4897.
[25] M. Ertürk, V. Karakaya, n-tuplet fixed point theorems for contractive type mappings in partially ordered metric spaces, Journal of Inequalities and Applications. 2013 (2013),196.
[26] A. Aghajani, R. Allahyari, M. Mursaleen, A generalization of Darbo's theorem with application to the solvability of systems of integral equations, Journal of Comp. and App. Math. 260 (2014), 68-77.
[27] V. Karakaya, N. E. H. Bouzara, K. Doğan and Y. Atalan, Existence of tripled fixed points for a class of condensing operators in Banach spaces, The Scientific World Journal. 2014 (2014), Article ID 541862.
[28] B. Samet, C. Vetro, Coupled fixed point theorems for multivalued nonlinear contraction mappings in partially ordered metric spaces, Nonlinear Anal. Theory Methods Appl. 74 (12) (2011), 4260-4268.
[29] K. P. R. Rao, G. N. V. Kishore, T. Kenan, A unique common tripled fixed point theorem for hybrid pair of maps, Abstract and Applied Analysis. 2012, Article ID 750403.
[30] B. C. Dhage, Some generalization of multi-valued version of Schauder's fixed point theorem with applications, CUBO A Mathematical J. 12 (2010), 139-151.
[31] R. Agarwal, M. Meehan, D. O'Regan, Fixed point theory and applications, Cambridge University, Cambridge, (2004).
[32] B. C. Dhage, Asymptotic stability of nonlinear functional integral equations via measure of noncompactness, Comm. Appl. Nonlinear Anal. 15(2) (2008), 89-101.
[33] S. Itoh, W. Takahashi, The common fixed point theory of singlevalued mappings and multivalued mappings, Pasific Journal of Mathematics. 79(2) (1978), 493-508.
[34] H. Thiems, Integrated semigroup and integral solutions to abstract Cauchy problem, J. Math. Anal. Appl. 152 (1990), 416-447.
[35] M. Kamenskii, V. Obukhovskii, P. Zecca, Condensing multivalued maps and semilinear differential inclusions in Banach spaces, Walter de Gruyter, Berlin-New York, (2001).
[36] R. R. Akmerov, M. I. Kamenski, A. S. Potapov, A. E. Rodkina and B. N. Sadovskii, Measure of noncompactness and condensing operators, Springer Basel, Birkhauser-Verlag, (1992).
[37] N. U. Ahmed, Semigroup theory with applications to systems and control, Harlow John Wiley & Sons Inc, New York, (1991).
[38] K. J. Engel, R. Nagel, One-parameter semigroups for linear evolution equations, Springer-Verlag, New York, (2000).
[39] A. Pazy, Semigroups of linear operators and applications to partial differential equations, Springer-Verlag, New York, (1983).
[40] C. Castaing, M. Valadier, Convex analysis and measurable multifunctions, Lecture Notes in Math. 580, Springer-Verlag, New York, (1977).

Department of Mathematics, Faculty of Arts and Sciences, Ahi Evran University, 40100 Kirşehir, Turkey
E-mail address: deryasekman@gmail.com

Faculty of Mathematics, University of Science and Technology Houari Boumediène, Bab-Ezzouar, 16111 Algies, Algeria
E-mail address: bzr.nour@gmail.com

Department of Mathematical Engineering, Yildiz Technical University, 34210 Istanbul, Turkey
E-mail address: vkkaya@yahoo.com